

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

ZIGLIP® 145 mg tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 145 mg fenofibrate (nanoparticles).
Excipients with known effect: Lactose anhydrous 444.1 mg.
For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Tablet.

White to off-white biconvex oblong tablets, debossed with "F" on one side and "145" on the other side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Ziglip 145 mg is indicated as an adjunct to diet and other non-pharmacological treatment (e.g. exercise, weight reduction) for the following:

- Treatment of severe hypertriglyceridaemia with or without low HDL cholesterol.
- Mixed hyperlipidaemia when a statin is contraindicated or not tolerated.
- Mixed hyperlipidaemia in patients at high cardiovascular risk in addition to a statin when triglycerides and HDL cholesterol are not adequately controlled.

4.2 Posology and method of administration

Dietary measures initiated before therapy should be continued. Response to therapy should be monitored by determination of serum lipid values. If an adequate response has not been achieved after several months (e.g. 3 months), complementary or different therapeutic measures should be considered.

APPROVED BY SCOMTE *1st*

EXPERT/DATA_ *Conf 13.02.11*

APPLICANT/DATA_ *P. Djogaliev 14.02.11*

Posology:

Adults:

The recommended dose is one tablet containing 145 mg fenofibrate taken once daily.

Special populations

Geriatric population:

In elderly patients, without renal impairment, the usual adult dose is recommended.

Renal impairment:

Dosage reduction is required in patients with renal impairment. In mild to moderate chronic kidney disease, lower doses of fenofibrate should be administered. In patients with severe chronic kidney disease, fenofibrate is not recommended.

Hepatic impairment:

Ziglip 145 mg is not recommended for use in patients with hepatic impairment due to the lack of data.

Paediatric population:

The safety and efficacy of fenofibrate in children and adolescents younger than 18 years has not been established. No data are available. Therefore the use of fenofibrate is not recommended in paediatric subjects under 18 years.

Method of administration:

Tablet may be given at any time of the day, with or without food (see section 5.2 Pharmacokinetic properties). Tablets should be swallowed whole with a glass of water.

4.3 Contraindications

- Hepatic insufficiency (including biliary cirrhosis and unexplained persistent liver function abnormality),
- Known gallbladder disease,
- Severe chronic kidney disease,
- Chronic or acute pancreatitis with the exception of acute pancreatitis due to severe hypertriglyceridemia,
- Known photoallergy or phototoxic reaction during treatment with fibrates or ketoprofen,
- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

APPROVED BY SCOMT/ISC²
EXPERT/DATA_ Prof 13.02.18
APPLICANT/DATA_ P. Sinyavski 14.02.18

4.4 Special warnings and precautions for use

Secondary causes of hyperlipidemia:

Secondary cause of hypercholesterolemia, such as uncontrolled type 2 diabetes mellitus, hypothyroidism, nephrotic syndrome, dysproteinemia, obstructive liver disease or alcoholism should be adequately treated before fenofibrate therapy is considered. Secondary cause of hypercholesterolemia related to pharmacological treatment can be seen with diuretics, β -blocking agents, estrogens, progestogens, combined oral contraceptives, immunosuppressive agents and protease inhibitors. In these cases it should be ascertained whether the hyperlipidaemia is of primary or secondary nature (possible elevation of lipid values caused by these therapeutic agents).

Liver function:

As with other lipid lowering agents, increases have been reported in transaminase levels in some patients. In the majority of cases these elevations were transient, minor and asymptomatic. It is recommended that transaminase levels are monitored every 3 months during the first 12 months of treatment and thereafter periodically. Attention should be paid to patients who develop increase in transaminase levels and therapy should be discontinued if AST (SGOT) and ALT (SGPT) levels increase to more than 3 times the upper limit of the normal range. When symptoms indicative of hepatitis occur (e.g. jaundice, pruritus), and diagnosis is confirmed by laboratory testing, fenofibrate therapy should be discontinued.

Pancreas:

Pancreatitis has been reported in patients taking fenofibrate (see sections Contraindications and Undesirable effects). This occurrence may represent a failure of efficacy in patients with severe hypertriglyceridemia, a direct drug effect, or a secondary phenomenon mediated through biliary tract stone or sludge formation with obstruction of the common bile duct.

Muscle:

Muscle toxicity, including rare cases of rhabdomyolysis, with or without renal failure, has been reported with administration of fibrates and other lipid-lowering agents. The incidence of this disorder increases in case of hypoalbuminaemia and previous renal insufficiency. Patients with pre-disposing factors for myopathy and/or rhabdomyolysis, including age above 70 years, personal or familial history of hereditary muscular disorders, renal impairment, hypothyroidism and high alcohol intake, may be at an increased risk of developing rhabdomyolysis. For these patients, the putative benefits and risks of fenofibrate therapy should be carefully weighed up. Muscle toxicity should be suspected in patients presenting diffuse myalgia, myositis, muscular cramps and weakness and/or marked increases in CPK (levels exceeding 5 times the upper normal range). In such cases treatment with fenofibrate should be stopped.

APPROVED BY SCOMTESSC

EXPERT/DATA_

APPLICANT/DATA_

13.02.18
N. Jovanović
14.02.18

The risk of muscle toxicity may be increased if the drug is administered with another fibrate or an HMG-CoA reductase inhibitor, especially in case of pre-existing muscular disease. Consequently, the co-prescription of fenofibrate with HMG-CoA reductase inhibitor or another fibrate should be reserved to patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and with a close monitoring of potential muscle toxicity.

Renal function:

Treatment should be interrupted in case of an increase in creatinine levels > 50% of ULN (upper limit of normal). It is recommended that creatinine is measured during the first 3 months after initiation of treatment and thereafter periodically (for dose recommendations, see section 4.2 Posology and method of administration).

Excipients:

As this medicinal product contains lactose, therefore patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Oral anticoagulants:

Fenofibrate enhances oral anticoagulant effect and may increase risk of bleeding. It is recommended that the dose of anticoagulants is reduced by about one third at the start of treatment and then gradually adjusted if necessary according to INR (International Normalised Ratio) monitoring.

Cyclosporin:

Some severe cases of reversible renal function impairment have been reported during concomitant administration of fenofibrate and cyclosporin. The renal function of these patients must therefore be closely monitored and the treatment with fenofibrate stopped in the case of severe alteration of laboratory parameters.

HMG-CoA reductase inhibitors and other fibrates:

The risk of serious muscle toxicity is increased if a fibrate is used concomitantly with HMG-CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution and patients monitored closely for signs of muscle toxicity (see section 4.4 Special warnings and precautions for use).

APPROVED BY SCDMT4 JSC
EXPERT/DATA - *Ref 13.02.18*
APPLICANT/DATA - *M. Sypucent*
14-02-18

Glitazones:

Some cases of reversible paradoxical reduction of HDL-cholesterol have been reported during concomitant administration of fenofibrate and glitazones. Therefore, it is recommended to monitor HDL-cholesterol if one of these components is added to the other and stopping of either therapy if HDL-cholesterol is too low.

Cytochrome P450 enzymes:

In vitro studies using human liver microsomes indicate that fenofibrate and fenofibric acid are not inhibitors of cytochrome (CYP) P450 isoforms CYP3A4, CYP2D6, CYP2E1, or CYP1A2. They are weak inhibitors of CYP2C19 and CYP2A6, and mild-to-moderate inhibitors of CYP2C9 at therapeutic concentrations.

Patients co-administered fenofibrate and CYP2C19, CYP2A6, and especially CYP2C9 metabolised drugs with a narrow therapeutic index should be carefully monitored and, if necessary, dose adjustment of these drugs is recommended.

4.6 Fertility, pregnancy and lactation

Pregnancy: There are no adequate data from the use of fenofibrate in pregnant women. Animal studies have not demonstrated any teratogenic effects. Embryotoxic effects have been shown at doses in the range of maternal toxicity (see section 5.3 Preclinical safety data). The potential risk for humans is unknown. Therefore, Ziglip 145 mg should only be used during pregnancy after a careful benefit/risk assessment.

Lactation: It is unknown whether fenofibrate and/or its metabolites are excreted in human milk. A risk to the suckling child cannot be excluded. Therefore fenofibrate should not be used during breast-feeding.

4.7 Effects on ability to drive and use machines

Ziglip 145 mg has no or negligible influence on the ability to drive and use machines.

4.8 Undesirable effects

The most commonly reported ADRs during fenofibrate therapy are digestive, gastric or intestinal disorders.

The following undesirable effects have been observed during placebo-controlled clinical trials (n=2344) and post-marketing with the below indicated frequencies:

MedDRA system organ class	Common ≥1/100, <1/10	Uncommon ≥1/1,000, <1/100	Rare ≥1/10,000, <1/1,000	Very rare <1/10,000	Frequency unknown (cannot be)

APPROVED BY SCOMT 5 JSC
 EXPERT/DATA - P. J. 15.02.18
 APPLICANT/DATA - P. J. 14.02.18

				0 incl. isolated reports	estimated from the available data)
Blood and lymphatic system disorders			Haemoglobin decreased White blood cell count decreased		
Immune system disorders			Hypersensitivity		
Nervous system disorders		Headache			
Vascular disorders		Thromboembolism (pulmonary embolism, deep vein thrombosis)*			
Respiratory, thoracic and mediastinal disorders					Interstitial lung disease ^a
Gastrointestinal disorders	Gastrointestinal signs and symptoms (abdominal pain, nausea, vomiting, diarrhoea, flatulence)	Pancreatitis*			
Hepatobiliary disorders	Transaminases increased (see section 4.4)	Cholelithiasis (see section 4.4)	Hepatitis		Jaundice, complications of cholelithiasis ^a (e.g. cholecystitis, cholangitis, biliary colic)

APPROVED BY SC/DHT/6/SC

EXPERT/DATA - Prof 13.02.18

APPLICANT/DATA - N. Sygarec
14.02.18

Skin and subcutaneous tissue disorders		Cutaneous hypersensitivity (e.g. rash, pruritus, urticaria)	Alopecia Photosensitivity reactions		Severe cutaneous reactions ^a (e.g. erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis)
Musculoskeletal, connective tissue and bone disorders		Muscle disorder (e.g. myalgia, myositis, muscular spasms and weakness)			Rhabdomyolysis ^a
Reproductive system and breast disorders		Sexual dysfunction			
Investigations		Blood creatinine increased	Blood urea increased		

*: In the FIELD-study, a randomized placebo-controlled trial performed in 9,795 patients with type 2 diabetes mellitus, a statistically significant increase in pancreatitis cases was observed in patients receiving fenofibrate versus patients receiving placebo (0.8% versus 0.5%; $p = 0.031$). In the same study, a statistically significant increase was reported in the incidence of pulmonary embolism (0.7% in the placebo group versus 1.1% in the fenofibrate group; $p = 0.022$) and a statistically non-significant increase in deep vein thromboses (placebo: 1.0% [48/4,900 patients] versus fenofibrate 1.4% [67/4,895 patients]; $p = 0.074$).

4.9 Overdose

Only anecdotal cases of fenofibrate overdosage have been received. In the majority of cases no overdose symptoms were reported.

No specific antidote is known. If an overdose is suspected, treat symptomatically and institute appropriate supportive measures as required. Fenofibrate cannot be eliminated by haemodialysis.

APPROVED BY SCDNTE/ISC

EXPERT/DATA - *Cep 12.02.18*

APPLICANT/DATA - *P. Jovanović*
14.02.18

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Serum Lipid Reducing Agents / Cholesterol and Triglycerides Reducers / Fibrates.

ATC code: C10AB05

Fenofibrate is a fibric acid derivative whose lipid modifying effects reported in humans are mediated via activation of Peroxisome Proliferator Activated Receptor type alpha (PPAR α).

Through activation of PPAR α , fenofibrate increases the lipolysis and elimination of atherogenic triglyceride-rich particles from plasma by activating lipoprotein lipase and reducing production of apoprotein CIII. Activation of PPAR α also induces an increase in the synthesis of apoproteins AI and AII.

The above stated effects of fenofibrate on lipoproteins lead to a reduction in very low- and low density fractions (VLDL and LDL) containing apoprotein B and an increase in the high density lipoprotein fraction (HDL) containing apoprotein AI and AII.

In addition, through modulation of the synthesis and the catabolism of VLDL fractions fenofibrate increases the LDL clearance and reduces small dense LDL, the levels of which are elevated in the atherogenic lipoprotein phenotype, a common disorder in patients at risk for coronary heart disease.

During clinical trials with fenofibrate, total cholesterol was reduced by 20 to 25%, triglycerides by 40 to 55% and HDL cholesterol was increased by 10 to 30%.

In hypercholesterolaemic patients, where LDL cholesterol levels are reduced by 20 to 35%, the overall effect on cholesterol results in a decrease in the ratios of total cholesterol to HDL cholesterol, LDL cholesterol to HDL cholesterol, or Apo B to Apo AI, all of which are markers of atherogenic risk.

There is evidence that treatment with fibrates may reduce coronary heart disease events but they have not been shown to decrease all cause mortality in the primary or secondary prevention of cardiovascular disease.

The Action to Control Cardiovascular Risk in Diabetes (ACCORD) lipid trial was a randomized placebo-controlled study of 5518 patients with type 2 diabetes mellitus treated with fenofibrate

APPROVED BY SCDMTE JSC

EXPERT/DATA_

APPLICANT/DATA_

Prof. B. 13.02.18
B. Djogacević
14.02.18

in addition to simvastatin. Fenofibrate plus simvastatin therapy did not show any significant differences compared to simvastatin monotherapy in the composite primary outcome of non-fatal myocardial infarction, non-fatal stroke, and cardiovascular death (hazard ratio [HR] 0.92, 95% CI 0.79-1.08, $p = 0.32$; absolute risk reduction: 0.74%). In the pre-specified subgroup of dyslipidaemic patients, defined as those in the lowest tertile of HDL-C (≤ 34 mg/dl or 0.88 mmol/L) and highest tertile of TG (≥ 204 mg/dl or 2.3 mmol/L) at baseline, fenofibrate plus simvastatin therapy demonstrated a 31% relative reduction compared to simvastatin monotherapy for the composite primary outcome (hazard ratio [HR] 0.69, 95% CI 0.49-0.97, $p = 0.03$; absolute risk reduction: 4.95%). Another prespecified subgroup analysis identified a statistically significant treatment-by-gender interaction ($p = 0.01$) indicating a possible treatment benefit of combination therapy in men ($p=0.037$) but a potentially higher risk for the primary outcome in women treated with combination therapy compared to simvastatin monotherapy ($p=0.069$). This was not observed in the aforementioned subgroup of patients with dyslipidaemia but there was also no clear evidence of benefit in dyslipidaemic women treated with fenofibrate plus simvastatin, and a possible harmful effect in this subgroup could not be excluded.

Extravascular deposits of cholesterol (tendinous and tuberous xanthoma) may be markedly reduced or even entirely eliminated during fenofibrate therapy.

Patients with raised levels of fibrinogen treated with fenofibrate have shown significant reductions in this parameter, as have those with raised levels of Lp(a). Other inflammatory markers such as C Reactive Protein are reduced with fenofibrate treatment.

The uricosuric effect of fenofibrate leading to reduction in uric acid levels of approximately 25% should be of additional benefit in those dyslipidaemic patients with hyperuricaemia.

Fenofibrate has been shown to possess an anti-aggregatory effect on platelets in animals and in a clinical study, which showed a reduction in platelet aggregation induced by ADP, arachidonic acid and epinephrine.

5.2 Pharmacokinetic properties

Ziglip 145 mg tablet contains 145 mg of fenofibrate nanoparticles.

Absorption:

Maximum plasma concentrations (C_{max}) occur within 2 to 4 hours after oral administration.

Plasma concentrations are stable during continuous treatment in any given individual.

Contrarily to previous fenofibrate formulations, the maximum plasma concentration and overall exposure of the nanoparticle formulation is independent from food intake. Therefore, Ziglip 145 mg may be taken without regard to meals.

APPROVED BY SCRM/JSC

EXPERT/DATA_

APPLICANT/DATA_

15.02.18

B. Gjorgjević
14.02.18

A food-effect study involving administration of the new 145 mg tablet formulation of fenofibrate to healthy male and female subjects under fasting conditions and with a high fat meal indicated that exposure (AUC and C_{max}) to fenofibric acid is not affected by food.

Distribution:

Fenofibric acid is strongly bound to plasma albumin (more than 99%).

Metabolism and excretion:

After oral administration, fenofibrate is rapidly hydrolysed by esterases to the active metabolite fenofibric acid. No unchanged fenofibrate can be detected in the plasma. Fenofibrate is not a substrate for CYP 3A4. No hepatic microsomal metabolism is involved.

The drug is excreted mainly in the urine. Practically all the drug is eliminated within 6 days. Fenofibrate is mainly excreted in the form of fenofibric acid and its glucuronide conjugate. In elderly patients, the fenofibric acid apparent total plasma clearance is not modified.

Kinetic studies following the administration of a single dose and continuous treatment have demonstrated that the drug does not accumulate. Fenofibric acid is not eliminated by haemodialysis.

The plasma elimination half-life of fenofibric acid is approximately 20 hours.

5.3 Preclinical safety data

Chronic toxicity studies have yielded no relevant information about specific toxicity of fenofibrate. Studies on mutagenicity of fenofibrate have been negative.

In rats and mice, liver tumours have been found at high dosages, which are attributable to peroxisome proliferation. These changes are specific to small rodents and have not been observed in other animal species. This is of no relevance to therapeutic use in man.

Studies in mice, rats and rabbits did not reveal any teratogenic effect. Embryotoxic effects were observed at doses in the range of maternal toxicity. Prolongation of the gestation period and difficulties during delivery were observed at high doses. No sign of any effect on fertility has been detected.

APPROVED BY SCOPJE JSC

EXPERT/DATA_ Bull 13.02.18

APPLICANT/DATA_ 10 G. Djurajević 14.02.18

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipient(s)

Lactose anhydrous
Hypromellose
Sodium laurilsulfate
Simethicone emulsion 30%
Croscarmellose sodium
Magnesium stearate

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

The tablets are immediate packed in PVC/PVdC/Al blisters, each blister containing 10 tablets.
The cardboard box contains three (3) blisters and a leaflet inside.

6.6 Special precautions for disposal and other handling

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

APPROVED BY SCOMTE JSC

EXPERT/DATA

APPLICANT/DATA

11

Handwritten signatures and dates:
13.02.18
14.02.18

Ziglip® 145 mg tablets

Summary of Product Characteristics



7. MARKETING AUTHORISATION HOLDER

ALKALOID AD Skopje
Blvd. Aleksandar Makedonski 12,
1000 Skopje, Republic of Macedonia
tel.: + 389 2 31 04 000
fax: +389 2 31 04 021
www.alkaloid.com.mk

8. MARKETING AUTHORISATION NUMBER(S)

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

10. DATE OF REVISION OF THE TEXT

APPROVED BY SCDA TE JSC

EXPERT/DATA

Conf 13.02.18

APPLICANT/DATA

P. Gjorgjievski
14.02.18