

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Influcid

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

1 tablet (250 mg) contains:

Active substances:

Aconitum trit. D3	25.0 mg
Gelsemium trit. D3	25.0 mg
Ipecacuanha trit. D3	25.0 mg
Phosphorus trit. D5	25.0 mg
Bryonia trit. D2	25.0 mg
Eupatorium perfoliatum trit. D1	25.0 mg

Excipients with known effect: lactose monohydrate, wheat starch

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Tablet

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Treatment and prevention of feverish flu-like infections and other acute viral upper respiratory tract infections. Supportive treatment of influenza.

4.2 Posology and method of administration

Posology

Adults and adolescents from 12 years of age in acute conditions, should take in 1 tablet every hour (but not more than 6 tablets per day). This frequency after a week of use can only be continued after consultation with a homeopathic experienced therapist. *With further improvement*, adults and adolescents over 12 years old should take 1 tablet 1-3 times a day.

Paediatric population

Children aged 1 to 6 years: in acute conditions, should not receive more than half of adult dose (3 tablets per day). For small children under 6 years of age, it is advisable to resolve the tablet before taking into a small amount of water.

Children aged 6-12 years should not receive more than 2/3 of adult dose (4 tablets per day).

Method of administration

Tablets should be allowed to dissolve slowly in the mouth. For young children, tablets can be dissolved in a little bit of water.

An interval of at least half an hour to meals should be kept.

4.3 Contraindications

Hypersensitivity to *Eupatorium perfoliatum* (common boneset) or other members of the asteraceae/compositae family, any other of the active substances or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

This medicinal product contains lactose and wheat starch. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicinal product. Wheat starch may contain gluten, but only in trace amounts, and is therefore considered safe for people with coeliac disease. Patients with wheat allergy (different from coeliac disease) should not take this medicinal product.

A reassessment of the therapy is necessary

- if condition does not improve or worsens,
- if additional complaints arise,
- if fever persists or rises above 39°C.

Paediatric population

Influcid should not be used in children under 1 year of age since its use is not sufficiently documented in this age group.

4.5 Interaction with other medicinal products and other forms of interaction

Interactions with other medicinal products have not been observed. The use of this homeopathic medicinal product does not exclude the use of other medicinal products.

Patients are advised to consult a doctor if they are taking, have recently taken or might take any other medicinal products.

Note: The action of homeopathic medicinal products may be adversely affected by generally negative factors of living habits as well as by stimulants or drink and tobacco.

4.6 Pregnancy and lactation

Patients are advised to consult a doctor before using this medicinal product.

Like all pharmaceuticals, Influcid must be taken during pregnancy and lactation only under consideration of the benefit-risk ratio assessment.

4.7 Effects on ability to drive and use machines

Influcid has no influence on the ability to drive and use machines.

4.8 Undesirable effects

Very rare (<1/10,000) or not known (frequency cannot be estimated from the available data):
Hypersensitivity reactions, e.g. skin rash and gastrointestinal complaints. In these cases, Influcid should be discontinued and appropriate treatment measures have to be taken.

Note: Existing complaints may temporarily become worse when homeopathic medicines are taken (initial homeopathic aggravation). In such cases, patients are advised to discontinue the medication and consult a doctor.

4.9 Overdose

No case of overdose has been reported.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Homeopathic medicinal product.

Homeopathic therapy differs markedly from conventional medicine with regard to the understanding of disease and the use of medicinal substances. Homeopathic remedies stimulate the organism's own defence mechanisms and thereby activate its self-healing power. The response of the organism to a homeopathic remedy is highly individual. For these reasons dose-specific pharmacological effects, pharmacodynamic or pharmacokinetic data cannot be determined.

5.1 Pharmacodynamic properties

Not applicable.

5.2 Pharmacokinetic properties

Not applicable.

5.3 Preclinical safety data

Not applicable. Influcid as a homeopathic pharmaceutical has not gone through preclinical testing.

6. PHARMACEUTICAL PROPERTIES

6.1 List of excipients

Lactose monohydrate, magnesium stearate, wheat starch.

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

5 years.

Do not use after the expiration date indicated on the package.

6.4 Special precautions for storage

Store at a temperature not exceeding 25 ° C. Store out of reach and sight of children.

6.5 Nature and contents of container

20 tablets in blister. Package containing 2 blisters and PIL.

6.6 Special precautions for disposal

No special requirements.

7. MANUFACTURER

Dr. Gustav Klein GmbH & Co. KG
Steinenfeld 3, 77736 Zell am Harmersbach, Germany

8. MARKETING AUTHORISATION HOLDER

Alpen Pharma GmbH, Steinenfeld 3,
77736 Zell am Harmersbach, Germany
Telephone: +49 (0) 7243 200 49 10
www.alpenpharma.com

9. MARKETING AUTHORISATION NUMBER(S)

[To be completed]

10. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

12/02/1997

11. DATE OF REVISION OF THE TEXT

May 2024